


muco^{derm}[®]

**FOR SOFT TISSUE THICKENING
IN DENTAL IMPLANTOLOGY**

soft tissue



WHY SOFT TISSUE AUGMENTATION:

Thin soft tissue (<2 mm) increases the risk of crestal bone loss^{1,2}, affects the aesthetic appearance, and impairs the dental hygiene³.

Augmentation of thin soft tissues helps to reduce alveolar bone loss and restore the hard and soft tissue profile^{2,4}.

mucoderm[®]

SOFT TISSUE AUGMENTATION AROUND TEETH AND IMPLANTS




mucoderm[®] is an acellular, dermal matrix of porcine origin indicated for the regeneration of soft tissue defects in dental, oral and maxillofacial surgery. Its native collagen structure serves as a scaffold for cells and blood vessels and is completely remodeled into patient's own soft tissue. The unique collagen architecture and exceptional biomechanical properties make mucoderm[®] a valid alternative to autologous connective tissue grafts in many indications.

1. Hämmerle C and Tarnow D 2018. J Periodontol.;89 Suppl 1:S291-S303.
2. Puišys A et al. 2015. Clin Oral Implants Res.;26(2):123-9.
3. Dhir S. 2011. J Indian Soc Periodontol. 15(2): 98-103.
4. Rossi AL et al. 2018. DentalCadmos ; 86(5):400-413.

WHY mucoderm[®]

FOR SOFT TISSUE AUGMENTATION

- + Significant increase of the (peri-implant) soft tissue in the horizontal and vertical dimension^{5,6,7}
- + Integrates well and gets completely remodeled^{6,7}
- + Less patient morbidity compared to autologous tissue transplants
- + Optimal thickness of 1.2-1.7mm
- + Excellent mechanical and volume stability even after hydration⁸
- + Easy to handle and to work with



5. Papi P et al. 2018. Biomed Res.;6406051.
6. Zafropoulos GG et al. 2016. J Indian Soc. Periodontol.; 20:441-5.
7. Puišys A et al. 2019. Int J Periodontics Restorative Dent.;39(3):341-347.
8. Kasaj A et al. 2016. Clin Oral Investig. 20(6):1303-15.

WHY SOFT TISSUE AUGMENTATION:

Thin soft tissue (<2 mm) increases the risk of crestal bone loss^{1,2}, affects the aesthetic appearance, and impairs the dental hygiene³.

Augmentation of thin soft tissues helps to reduce alveolar bone loss and restore the hard and soft tissue profile^{2,4}.

mucoderm®

SOFT TISSUE AUGMENTATION AROUND TEETH AND IMPLANTS

PERI-IMPLANT MUCOSAL THICKENING - 5 YEARS FOLLOW-UP

CLINICAL CASE BY

Dr. Algirdas Puišys, Vilnius, Lithuania



Crestal incision at the edentulous ridge and raising of a full-thickness flap buccally and lingually
 Hydrated mucoderm® punched and pulled over the healing cap
 The margins of the flap are adapted and sutured leaving the abutment open
 Situation after suture removal, one week post-operative
 Final restoration five months post-operative
 Stable clinical situation after five years

BUCCAL SOFT TISSUE AUGMENTATION AT SINGLE IMPLANT SITE⁹

CLINICAL CASE BY

Dr. Martina Stefanini and Prof. Dr. Giovanni Zucchelli, Bologna, Italy



Occlusal view of initial clinical situation of missing soft tissue at buccal side
 mucoderm® application following hydration in sterile saline
 mucoderm® placed to thicken the buccal soft tissue
 Flap closure
 Lateral situation at one year
 Occlusal view showing soft tissue increase at one year follow-up

„Increase soft tissue thickness to improve aesthetics and to promote healthy tissues is our objective, but reducing patient morbidity is of paramount importance: with mucoderm® you can reach both endpoints.“
Dr. Martina Stefanini

>100 scientific contributions
 demonstrating the unique properties and clinical success of mucoderm®

SOCKET SEALING WITH MUCODERM® FOR LATER IMPLANT PLACEMENT⁴

CLINICAL CASE BY

Dr. Alessandro Rossi, Milan, Italy



Initial clinical situation showing strongly compromised tooth 21
 Extraction socket filled with bone substitute material
 Application of mucoderm® to seal the socket and to thicken the soft tissue buccally
 Wound closure. 2/3 of mucoderm® are exposed
 Eight weeks after surgery
 Implant placement

- Hydration in sterile saline or blood prior to application
- After hydration easy size adaptation and punching
- For peri-implant soft tissue thickening mucoderm® can be placed in full-flap approach directly on the bone
- For peri-implant soft tissue thickening mucoderm® must be covered to ensure its revitalization; a minor surface area can be left exposed
- Open healing possible in socket preservation with best results regarding thickening if only 1/3 of the matrix is exposed

For more information on indications and clinical cases:
www.INDICATION-MATRIX.com
 or have a look in the mucoderm® **SURGICAL GUIDE!**



mucoderm® is an acellular, dermal matrix of porcine origin indicated for the regeneration of soft tissue defects in dental, oral and maxillofacial surgery. Its native collagen structure serves as a scaffold for cells and blood vessels and is completely remodeled into patient's own soft tissue. The unique collagen architecture and exceptional biomechanical properties make mucoderm® a valid alternative to autologous connective tissue grafts in many indications.

1. Hämmerle C and Tarnow D 2018. J Periodontol.;89 Suppl 1:S291-S303.
 2. Puišys A et al. 2015. Clin Oral Implants Res.;26(2):123-9.
 3. Dhir S. 2011. J Indian Soc Periodontol. 15(2): 98-103.
 4. Rossi AL et al. 2018. DentalCadmos ; 96(5):400-413.

9. Stefanini M, Randon A and Zucchelli G. 2019. Porcine-derived acellular dermal matrix for buccal soft tissue augmentation at implant sites: a 1 year follow-up case series. Int. J of Periodontics and Restorative Dentistry, in press

bone & tissue
regeneration

botiss
biomaterials

Innovation.
Regeneration.
Aesthetics.

soft tissue

botiss biomaterials GmbH
Hauptstr. 28
15806 Zossen / Germany

education

Tel.: +49 33769 / 88 41 985
Fax: +49 33769 / 88 41 986

hard tissue

www.botiss.com
www.botiss-dental.com
facebook: botissdental